Author Index

Abraham, M. H.
——, Buist, G. J., Grellier, P. L., McGill, R. A., Doherty, R. M. Kamlet, M. J., Taft, R. W. and Maroldo, S. G.
Solubility properties in polymers and biological media. II. A new method for the characterisation of the adsorption of gases and vapours on solids 15
Amor, J.-C., see Pedroso, E. 281
Armstrong, D. W., see Spino, L. A. 147
Ashoor, S. H.
—— and Knox, M. J.
Determination of vitamin A derivatives in fortified foods and commercial vitamin supplements by high-performance liquid chromatography 419
Aue, W. A., see Tang, Y.-Z. 125, 243
Beger, J., see Voelkel, A. 29
Behar, N., see Sébille, B. 61
Bidleman, T. F., see Zaranski, M. T. 235
Budevsky, O., see Dimitrova, B. 81
Buist, G. J., see Abraham, M. H. 15
Burger, G., see Pochard, M.-F. 315
Cachet, Th.
——, Kibwage, I. O., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.
Optimization of the separation of erythromycin and related substances by high-performance liquid chromatography 91
Cazau, M., see Oliveros, L. 183
Chang, C. A., see Lin, G.-C. 371
Cheng, Y.-H., see Sagara, K. 365
Chenieux, J. C., see Montagu, M. 426
Chevalier, M., see Pochard, M.-F. 315
Dahiya, J.S.
Reversed-phase high-performance liquid chromatography of *Cajanus cajan* phytoalexins 355
Dimitrova, B.
—— and Budevsky, O.
Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by high-performance liquid chromatography 81
Doherty, R. M., see Abraham, M. H. 15
Ebert, K., see Voelkel, A. 29
Frey, D. D.
Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography 1
Fritz, J. S., see Tanaka, K. 271
Fulton, S., see Kopaciewicz, W. 111
Galli, B.
——, Gasparrini, F., Misiti, D., Natile, G. and Palmieri, G.
Thin-layer chromatography of metal complexes on glycidoxypropyl functionalized silica plates 377
Gasparrini, F., see Galli, B. 377
Gattrell, M. A.
—— and Kirk, D. W.
Interconversion of hydroquinone and benzoquinone during styrene–divinylbenzene resin-based ion-exclusion chromatography 404
Gercken, G., see Schatowitz, B. 43
Gimenez-Gallego, G.
—— and Thomas, K. A.
High-performance liquid chromatography of phenylthiocarbamyl-amino acids. Application to carboxyl-terminal sequencing of proteins 299
Giralt, E., see Pedroso, E. 281
Gleizes, E., see Pochard, M.-F. 315
Grandas, A., see Pedroso, E. 281
Greenaway, W.
—— and Whatley, F. R.
Formation of N-carboxy compounds during trimethylsilanization of amino acids in the presence of carbon dioxide 383
Grellier, P. L., see Abraham, M. H. 15
Griest, W. H., see Row, K. H. 193
Grimsrud, E. P., see Jones, C. S. 139
Hansen, S. H.
——, Helboe, P. and Thomsen, M.
Separation of basic drug substances by reversed-phase high-performance liquid chromatography on dynamically modified silica and on bonded-phase materials 71
Hartwick, R. A., see Raglione, T. V. 205, 213
Helboe, P., see Hansen, S. H. 71
Hirayama, C.
——, Ihara, H., Shiba, M., Nakamura, M., Motozato, Y. and Kunitake, T.
Macroporous glucomannan spheres for the size-exclusion separation of aqueous macromolecules 175
Hizukuri, S., see Koizumi, K. 396
Hodges, R. S., see Mant, C. T. 155
Hoogmartens, J., see Cachet, Th. 91
Hurst, W. J.
——, Snyder, K. P. and Martin, Jr., R. A.
Determination of aflatoxins in peanut products using disposable bonded-phase columns and post-column reaction detection 413

Author Index

Abraham, M. H.
——, Buist, G. J., Grellier, P. L., McGill, R. A., Doherty, R. M. Kamlet, M. J., Taft, R. W. and Maroldo, S. G.
Solubility properties in polymers and biological media. II. A new method for the characterisation of the adsorption of gases and vapours on solids 15
Amor, J.-C., see Pedroso, E. 281
Armstrong, D. W., see Spino, L. A. 147
Ashoor, S. H.
—— and Knox, M. J.
Determination of vitamin A derivatives in fortified foods and commercial vitamin supplements by high-performance liquid chromatography 419
Aue, W. A., see Tang, Y.-Z. 125, 243
Beger, J., see Voelkel, A. 29
Behar, N., see Sébille, B. 61
Bidleman, T. F., see Zaranski, M. T. 235
Budevsky, O., see Dimitrova, B. 81
Buist, G. J., see Abraham, M. H. 15
Burger, G., see Pochard, M.-F. 315
Cachet, Th.
——, Kibwage, I. O., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.
Optimization of the separation of erythromycin and related substances by high-performance liquid chromatography 91
Cazau, M., see Oliveros, L. 183
Chang, C. A., see Lin, G.-C. 371
Cheng, Y.-H., see Sagara, K. 365
Chenieux, J. C., see Montagu, M. 426
Chevalier, M., see Pochard, M.-F. 315
Dahiya, J.S.
Reversed-phase high-performance liquid chromatography of *Cajanus cajan* phytoalexins 355
Dimitrova, B.
—— and Budevsky, O.
Investigation of an alkylamine modified and pH-controlled mobile phase for separation of pyrazolone derivatives by high-performance liquid chromatography 81
Doherty, R. M., see Abraham, M. H. 15
Ebert, K., see Voelkel, A. 29
Frey, D. D.
Free energy consumption at self-sharpening concentration fronts in isothermal fixed-bed adsorption. Theory for isotherms of arbitrary shape and applications in displacement chromatography 1
Fritz, J. S., see Tanaka, K. 271
Fulton, S., see Kopaciewicz, W. 111
Galli, B.
——, Gasparrini, F., Misiti, D., Natile, G. and Palmieri, G.
Thin-layer chromatography of metal complexes on glycidoxypropyl functionalized silica plates 377
Gasparrini, F., see Galli, B. 377
Gattrell, M. A.
—— and Kirk, D. W.
Interconversion of hydroquinone and benzoquinone during styrene–divinylbenzene resin-based ion-exclusion chromatography 404
Gercken, G., see Schatowitz, B. 43
Gimenez-Gallego, G.
—— and Thomas, K. A.
High-performance liquid chromatography of phenylthiocarbamyl-amino acids. Application to carboxyl-terminal sequencing of proteins 299
Giralt, E., see Pedroso, E. 281
Gleizes, E., see Pochard, M.-F. 315
Grandas, A., see Pedroso, E. 281
Greenaway, W.
—— and Whatley, F. R.
Formation of N-carboxy compounds during trimethylsilanization of amino acids in the presence of carbon dioxide 383
Grellier, P. L., see Abraham, M. H. 15
Griest, W. H., see Row, K. H. 193
Grimsrud, E. P., see Jones, C. S. 139
Hansen, S. H.
——, Helboe, P. and Thomsen, M.
Separation of basic drug substances by reversed-phase high-performance liquid chromatography on dynamically modified silica and on bonded-phase materials 71
Hartwick, R. A., see Raglione, T. V. 205, 213
Helboe, P., see Hansen, S. H. 71
Hirayama, C.
——, Ihara, H., Shiba, M., Nakamura, M., Motozato, Y. and Kunitake, T.
Macroporous glucomannan spheres for the size-exclusion separation of aqueous macromolecules 175
Hizukuri, S., see Koizumi, K. 396
Hodges, R. S., see Mant, C. T. 155
Hoogmartens, J., see Cachet, Th. 91
Hurst, W. J.
——, Snyder, K. P. and Martin, Jr., R. A.
Determination of aflatoxins in peanut products using disposable bonded-phase columns and post-column reaction detection 413

Ihara, H., see Hirayama, C. 175
Johnsen, P. B.
—— and Kuan, J.-C. W.
Simplified method to quantify geosmin and 2-methylisoborneol concentrations in water and microbiological cultures 337
Jones, C. S.
—— and Grimsrud, E. P.
Mass spectrometric measurements of the negative ions formed in the thermionic ionization detector with nitrogen carrier gas 139
Kamlet, M. J., see Abraham, M. H. 15
Keller, M. A.
—— and Saba, C. S.
Chromatographic analysis of phenyl-1-naphthylamine and 4,4′-dioctyldiphenylamine and their intermediate oxidation products in oxidized lubricants 325
Kennerly, D. A.
Molecular species analysis of lysophospholipids using high-performance liquid chromatography and argentation thin-layer chromatography 291
Kibwage, I. O., see Cachet, Th. 91
Kinjo, S., see Ohta, H. 409
Kirk, D. W., see Cattrell, M. A. 404
Knox, M. J., see Ashoor, S. H. 419
Koizumi, K.
——, Utamura, T., Kubota, Y. and Hizukuri, S.
Two high-performance liquid chromatographic columns for analyses of malto-oligosaccharides 396
Kopaciewicz, W.
——, Fulton, S. and Lee, S. Y.
Influence of pore and particle size on the frontal uptake of proteins. Implications for preparative anion-exchange chromatography 111
Krajewska, A. M.
—— and Powers, J. J.
Gas chromatography of methyl derivatives of naturally occurring capsaicinoids 223
Kuan, J.-C. W. see Johnsen, P. B. 337
Kubota, Y., see Koizumi, K. 396
Kunitake, T., see Hirayama, C. 175
Leem S. Y., see Kopaciewicz, W. 111
Levillain, P., see Montagu, M. 426
Lin, G.-C.
—— and Chang, C. A.
Heavy alkylated benzene stationary phase for the gas–liquid chromatographic separation of ethylbenzene, xylenes, styrene, iso- and *n*-propylbenzene 371
Lochmüller, C. H.
—— and Mink, L. P.
Adsorption isotherms of ethyl acetate modifier on silica from supercritical carbon dioxide 55
Locke, D. C., see Schneiderman, M. A. 343.
McGill, R. A., see Abraham, M. H. 15
Mant, C. T.
—— and Hodges, R. S.
On-line derivatization of silica supports for regeneration and preparation of reversed-phase columns 155
Mareš, P., see Řezanka, T. 390
Maroldo, S. G., see Abraham, M. H. 15
Martin, Jr., R. A., see Hurst, W. J. 413
Maskarinec, M. P., see Row, K. H. 193
Mink, L. P., see Lochmüller, C. H. 55
Misiti, D., see Galli, B. 377
Montagu, M.
——, Levillain, P., Chenieux, J. C. and Rideau, M.
Fluorodensitometric assay of 6*H*-pyrido[4,3-*b*]-5,11-dimethylcarbazoles (ellipticine and derivatives) biosynthesized by *Ochrosia elliptica* cultures *in vitro* 426
Motozato, Y., see Hirayama, C. 175
Nakamura, M., see Hirayama, C. 175
Natile, G., see Galli, B. 377
Nishikawa, T., see Onodera, S. 259
Nomura, M., see Yamamoto, S. 101
Ohta, H.
——, Kinjo, S. and Osajima, Y.
Glass capillary gas chromatographic analysis of volatile components of canned Philippine pineapple juice 409
Oliveros, L.
—— and Cazau, M.
Étude de la résolution d'une série d'oxydes de phosphines racémiques par chromatographie en phase liquide sur deux phases stationnaires chirales de même nature et de structures voisines 183
Onodera, S.
——, Nishikawa, T. and Suzuki, S.
Chemical changes of organic compounds in chlorinated water. XIV. Characterization and determination of halogenated organics formed during chlorination of water from the Tama river 259
Osajima, Y., see Ohta, H. 409
Oshima, T., see Sagara, K. 365
Palmieri, G., see Galli, B. 377
Pedroso, E.
——, Grandas, A., Amor, J.-C. and Giralt, E.
Reversed-phase high-performance liquid chromatography of protected peptide segments 281
Piquion, J., see Sébille, B. 61
Pochard, M.-F.
——, Burger, G., Chevalier, M. and Gleizes, E.
Détermination des résidus de chloramphénicol par chromatographie liquide haute performance en phase inverse. Application à une étude pharmacocinétique chez la truite arc-en-ciel avec confirmation par spectrométrie de masse 315

Powers, J. J., see Krajewska, A. M. 223
Raglione, T. V.
——, Troskosky, J. A. and Hartwick, R. A.
On-line microbore high-performance liquid chromatography–capillary gas chromatography–mass spectrometry. II. Application to the analysis of solvent refined coal 213
——, Troskosky, J. A. and Hartwick, R. A.
On-line microbore high-performance liquid chromatography–capillary gas chromatography–mass spectrometry. I. Development of isotachic eluent splitters 205
Rideau, M., see Montagu, M. 426
Regnier, F. E., see Stringham, R. W. 305
Řezanka, T.
—— and Mareš, P.
Unusual and very long-chain fatty acids produced by Basidiomycetes 390
Roets, E., see Cachet, Th. 91
Roth, M.
Effect of column inlet concentration profile on the sensitivity of analysis. A quantitative assessment 360
Row, K. H.
——, Griest, W. H. and Maskarinec, M. P.
Separation of modified nucleic acid constituents by micellar electrokinetic capillary chromatography 193
Saba, C. S., see Keller, M. A. 325
Sagara, K.
——, Oshima, T., Yoshida, T., Ton, Y.-Y., Zhang, G. and Chen, Y.-H.
Determination of Cinnamomi Cortex by high-performance liquid chromatography 365
Sano, Y., see Yamamoto, S. 101
Schatowitz, B.
—— and Gercken, G.
Benzyl esters of C_2–C_{20} fatty acids and metabolically relevant carboxylic acids. Preparation and gas chromatography–mass spectrometry 43
Schneiderman, M. A.
——, Sharma, A. K. and Locke, D. C.
Determination of anthraquinone in paper and wood using supercritical fluid extraction and high-performance liquid chromatography with electrochemical detection 343
Sébille, B.
——, Thuaud, N., Piquion, J. and Behar, N.
Determination of association constants of β-cyclodextrin and β-cyclodextrin-bearing polymers with drugs by high-performance liquid chromatography 61
Sharma, A. K., see Schneiderman, M. A. 343
Shiba, M., see Hirayama, C. 175
Snyder, K. P., see Hurst, W. J. 413
Spino, L. A.
——, Armstrong, D. W. and Vo-Dinh, T.
Direct synchronous luminescence detection of co-eluting solutes in pseudophase liquid chromatography 147
Stringham, R. W.
—— and Regnier, F. E.
Selective non-adsorption preparative chromatography of bovine immunoglobulin G1 305
Suzuki, S., see Onodera, S. 259
Szymanowski, J., see Voelkel, A. 29
Taft, R. W., see Abraham, M. H. 15
Tanaka, K.
—— and Fritz, J. S.
Ion-exclusion chromatography of non-ionic substances with conductivity detection 271
Tang, Y.-Z.
—— and Aue, W. A.
Gas chromatographic determination of benzaldehyde by gas-phase luminescence 243
——, and Aue, W. A.
General photometric detector for gas chromatography 125
Thomas, K. A., see Gimenez-Gallego, G. 299
Thomsen, M., see Hansen, S. H. 71
Thuaud, N., see Sébille, B. 61
Tong, Y.-Y., see Sagara, K. 365
Troskosky, J. A., see Raglione, T. V. 205, 213
Utamura, T., see Koizumi, K. 396
Vanderhaeghe, H., see Cachet, Th. 91
Vo-Dinh, T., see Spino, L. A. 147
Voelkel, A.
——, Szymanowski, J., Beger, J. and Ebert, K.
Polarity of 1,3-bis[ω-alkoxyoligo(oxyethylene)]propan-2-ols as measured by reversed-phase gas chromatography 29
Whatley, F. R., see Greenaway, W. 383
Yamamoto, S.
——, Nomura, M. and Sano, Y.
Resolution of proteins in linear gradient elution ion-exchange and hydrophobic interaction chromatography 101
Yasuhara, A.
Comparison of volatile components between fresh and rotten mussels by gas chromatography–mass spectrometry 251
Yoshida, T., see Sagara, K. 365
Zaranski, M. T.
—— and Bidleman, T. F.
High-volume elution chromatography of dichlorobenzenes on a polyurethane foam–Tenax sandwich cartridge 235
Zhang, G., see Sagara, K. 365